

Dynamics of Peakons, Jetlets and G-strands

Darryl D Holm, Imperial College London, UK, *d.holm@ic.ac.uk

Abstract

The way we view hydrodynamics changed forever when Arnold made his revolutionary discovery [1] that the Euler equations for an ideal fluid represent geodesic motion on SDiff (volume preserving diffeomorphisms) with respect to the L^2 norm on the tangent space $\text{TSDiff} \simeq \mathfrak{X}_{div} \times \text{SDiff}$, where \mathfrak{X}_{div} denotes the divergence-free vector fields. Arnold's famous paper has led to many further developments in continuum dynamics. These developments range, for example, from shallow-water solitons to shape analysis for computational anatomy.

The developments of Arnold's discovery that we will discuss in this talk are based on a dual pair of momentum maps that emerge from the Euler-Poincaré theory of Lagrangian reduction by symmetry when the symmetry is the Lie group of diffeomorphisms acting on a smooth manifold M , or on a space of smooth embeddings in M [2].

The examples we shall discuss as variations on the theme of dual momentum maps are:

1. Shallow-water solitons called peakons
2. Jetlets: a new type of coherent particle-like fluid excitation that carries momentum and angular momentum, while preserving its circulation.
3. G-strands: maps from \mathbb{R}^2 (or \mathbb{C}) into a Lie group G that are determined from Hamilton's principle for a Lagrangian that is *invariant* under G . For continuum dynamics in a domain M , the group is $G = \text{Diff}(M)$.

If time remains, we will also say a word about stochastic extensions of these examples.

References

- [1] Arnold, V. I., "Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits," *Annales de l'institut Fourier*, **6**, No. 1, 319–361 (1966).
- [2] Holm, D. D., Marsden, J. E., "Momentum Maps and Measure-valued Solutions," in: *The Breadth of Symplectic and Poisson Geometry*, J.E. Marsden and T.S. Ratiu, Editors, Birkhäuser Boston, Boston, MA, 2004, *Progr. Math.*, **232**, pp. 203–235.