

NUMERICAL MINIMIZATION OF DIRICHLET LAPLACIAN EIGENVALUES OF FOUR-DIMENSIONAL GEOMETRIES*

PEDRO R. S. ANTUNES[†] AND ÉDOUARD OUDET[‡]

Abstract. We develop the first numerical study in four dimensions of optimal eigenmodes associated with the Dirichlet Laplacian. We describe an extension of the method of fundamental solutions adapted to the four-dimensional context. Based on our numerical simulation and a postprocessing adapted to the identification of relevant symmetries, we provide and discuss the numerical description of the eighth first optimal domains.

Key words. Dirichlet Laplacian, eigenvalues, higher-dimensional space, shape optimization

AMS subject classifications. 35Q93, 65N25

DOI. 10.1137/16M1083773

1. Introduction. Shape and design optimization problems involving eigenvalues of elliptic operators is a very active topic in spectral theory and arises in many physical applications. A better understanding of this questions allows, for example, us to tune and improve the acoustic properties of musical instruments [8, 15, 25, 34, 36] or to find optimal designs of composite materials (e.g., [2]). A typical situation in engineering is the identification of shapes or designs of structures to prescribe their mechanical properties: being either as rigid or as soft as possible (e.g., [24]), avoiding torsional oscillations (e.g., [10]), etc. Such problems appear also in the context of electromagnetism when considering the design of optimal accelerator cavities (e.g., [1]).

A prototype of these shape optimization problems is the minimization of Dirichlet eigenvalues of the Laplacian, with a volume constraint. Whereas classical and widely studied, this problem collects all the main computational difficulties of design optimization: nonsmoothness, local minima, a high number of degrees of freedom, and a computationally expensive cost function.

Let $\Omega \subset \mathbb{R}^d$ be a bounded open set, not necessarily connected, and consider the Dirichlet eigenvalue problem

$$(1) \quad \begin{cases} -\Delta u = \lambda u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

defined in the Sobolev space $H_0^1(\Omega)$. We will denote the eigenvalues by $0 < \lambda_1(\Omega) \leq \lambda_2(\Omega) \leq \dots$ where each $\lambda_k(\Omega)$ is counted with its multiplicity and the corresponding orthonormal real eigenfunctions by u_i , $i = 1, 2, \dots$. The shape optimization can be formulated as determining

$$(2) \quad \lambda_k^* = \text{Min} \{ \lambda_k(\Omega) : \Omega \subset \mathbb{R}^d, |\Omega| = 1 \}, \quad k = 1, 2, \dots$$

*Submitted to the journal's Computational Methods in Science and Engineering section July 7, 2016; accepted for publication (in revised form) January 5, 2017; published electronically June 13, 2017.

<http://www.siam.org/journals/sisc/39-3/M108377.html>

Funding: The work of the first author was partially supported by FCT, Portugal, through the program “Investigador FCT” with reference IF/00177/2013 and the scientific project PTDC/MAT-CAL/4334/2014. The work of the second author was supported by the ANR, through the projects COMEDIC, PGMO, and OPTIFORM.

[†]Group of Mathematical Physics, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, Edifício C6 1749-016 Lisboa, Portugal (prantunes@fc.ul.pt).

[‡]LJK, Université de Grenoble, 51 rue des Mathématiques, F-38041 Grenoble Cedex 9 (edouard.oudet@imag.fr).

The existence of minimizing domains among quasi-open sets was recently proved (cf. [12, 31]) and some numerical studies allowed one to suggest candidates to be minimizers at the beginning of the spectrum and also to explore some properties such as connectedness, symmetry, and multiplicity of optimal eigenvalues (e.g., [35, 6, 37]).

In this paper we address the solution of shape optimization problems for the Dirichlet eigenvalues of the Laplacian of four-dimensional (4D) geometries. Besides the fact that differential geometry may present interesting features (e.g., [17, 21, 42]), the solution of 4D shape optimization problems and the calculation of eigenvalues of 4D geometries are also challenging from the computational point of view. Indeed, to the best of our knowledge this is the first time that a 4D shape optimization related to eigenvalues is considered. For this purpose we used the method of fundamental solutions (MFS) as solver for the eigenvalue problem. The MFS is a mesh-free method and thus, in this context, it avoids meshing 4D domains which could be a difficult and time consuming task. Moreover, the huge dimension of the matrices associated with classical mesh-type methods such as finite element methods would probably be prohibitive for solving the eigenvalue problem accurately in a reasonable time. On the other hand, the problem can be circumvented using spectral methods, such as the MFS. We can prove that the solution of the eigenvalue problem is also a solution of an integral equation defined on the boundary of a domain which can be seen as a *space dimension reduction* and the problem of solving an eigenvalue problem with 4D domain is replaced by the solution of an integral equation defined on a three-dimensional (3D) hypersurface.

The convergence of the MFS, when applied to smooth shapes, is known to be very fast, in some cases even exponential (cf. [27, 4, 9]), which ensures that this integral equation can be solved accurately with relatively small dimension matrices.

The plan of the paper is the following. In section 2 we give a brief description of some aspects of the numerical approach, namely, we define a parameterization of a general star-shaped domain in terms of 4D hyperspherical harmonics, we present an algorithm for the distribution of the collocation points and source points for the MFS, and use the Betcke–Trefethen subspace angle approach for the eigenvalue calculation and describe an algorithm for the optimization. Section 3 describes an algorithm for the visualization of the optimal shapes and presents some of the numerical results that we gathered. In section 4 we discuss some of the results that we obtained.

2. Numerical solution of the shape optimization problem.

2.1. Brief description of the MFS. We will use the MFS as forward solver for the eigenvalue problem. We take a fundamental solution of the Helmholtz equation in \mathbb{R}^d ,

$$(3) \quad \Phi_\lambda(x) = \frac{i}{4} \left(\frac{\sqrt{\lambda}}{2\pi|x|} \right)^{\frac{d-2}{d}} H_{\frac{d-2}{2}}^{(1)}(\sqrt{\lambda}|x|),$$

where $|\cdot|$ denotes the Euclidean norm in \mathbb{R}^d and $H_{\frac{d-2}{2}}^{(1)}$ is a Hankel function defined by Bessel functions (e.g., [32]),

$$H_{\frac{d-2}{2}}^{(1)}(\cdot) = J_{\frac{d-2}{2}}(\cdot) + iY_{\frac{d-2}{2}}(\cdot).$$

Now we introduce the following.

DEFINITION 2.1. *An admissible source set $\hat{\Gamma}$ is the boundary of a bounded open set $\hat{\Omega}$ such that $\hat{\Omega} \subset \Omega$ with $\hat{\Gamma}$ surrounding $\partial\Omega$.*

The MFS approximation is a linear combination

$$(4) \quad u \approx \tilde{u}(x) = \sum_{i=1}^M \beta_i \phi_j(x),$$

where

$$(5) \quad \phi_j = \Phi_\lambda(\cdot - y_j)$$

are M point sources centered at some points y_j that are placed on an admissible source set $\hat{\Gamma}$. By construction, the MFS approximation (4) satisfies the PDE of the eigenvalue problem (1) and the numerical accuracy of the solution is related to the accuracy of the approximation of the boundary condition. As proven in [5], given an admissible source set $\hat{\Gamma}$, we have the following result.

THEOREM 2.2. *If $\hat{\Gamma}$ is an admissible source set, then*

$$S(\hat{\Gamma}) = \text{span} \left\{ \Phi_\lambda(\cdot - y)|_\Omega : y \in \hat{\Gamma} \right\}$$

is dense in $\mathcal{H}_\lambda(\Omega) = \{v \in H^1(\Omega) : (\Delta + \lambda)v = 0\}$ with the $H^1(\Omega)$ topology.

By the Sobolev regularity theorem (cf. [26]), assuming that Ω is $C^{1,2}$, we know that the eigenfunctions belong to the Sobolev space $H^2(\Omega)$ and, thus, the density theorem Theorem 2.2 states that an eigenfunction can be arbitrarily well approximated by an MFS linear combination (4).

2.2. Parameterization of domains. We will restrict the shape optimization (2) to a similar problem defined in the class of sets that are finite unions of star-shaped domains. This restriction is only related to the simplicity of parameterizing a generic star-shaped domain and not to limitations of the MFS which can also be applied for nonsimply connected domains (cf. [13]). Moreover, using the Wolf–Keller theorem, if some optimizer for a given eigenvalue λ_k is a finite union of star-shaped domains, then each of its connected components is a minimizer for a previous eigenvalue. Thus, we can restrict our attention to star-shaped domains and study the disconnected case using the Wolf–Keller result ([43]).

A 4D star-shaped domain is isometric to the domain whose boundary is parameterized by

$$(6) \quad \begin{cases} x = r(\beta, \theta, \phi) \sin(\beta) \sin(\theta) \cos(\phi), \\ y = r(\beta, \theta, \phi) \sin(\beta) \sin(\theta) \sin(\phi), \\ z = r(\beta, \theta, \phi) \sin(\beta) \cos(\theta), \\ w = r(\beta, \theta, \phi) \cos(\beta), \end{cases}$$

where $r(\beta, \theta, \phi) > 0$ for $\beta \in [0, \pi]$, $\theta \in [0, \pi]$, and $\phi \in [0, 2\pi[$.

The Laplace–Beltrami operator on S^3 is defined by

$$\Delta_{S^3} = \frac{1}{\sin^2 \beta} \frac{\partial}{\partial \beta} \sin^2 \beta \frac{\partial}{\partial \beta} + \frac{1}{\sin^2 \beta} \Delta_{S^2},$$

where Δ_{S^2} is the Laplace–Beltrami operator on the unitary sphere S^2 . We define the family of 4D hyperspherical harmonics (4D HSH) by

$$(7) \quad \begin{aligned} n &= 0, 1, 2, \dots, & \beta &\in [0, \pi], \\ S_{nl}^m(\beta, \theta, \phi) &= c_{n,l,m} \sin^l(\beta) C_{n-l}^{l+1}(\cos(\beta)) Y_l^m(\theta, \phi)(\beta, \theta, \phi), & 0 \leq l \leq n, & \theta \in [0, \pi], \\ & & -l \leq m \leq l, & \phi \in [0, 2\pi[, \end{aligned}$$

where Y_l^m are 3D spherical harmonics, C_{n-l}^{l+1} are Gegenbauer polynomials, and

$$c_{n,l,m} = 2^{l+\frac{1}{2}} \sqrt{\frac{(n+1)\Gamma(n-l+1)}{\pi\Gamma(n+l+2)}} \Gamma(l+1).$$

The 4D HSH are eigenfunctions of Δ_{S^3} ,

$$\Delta_{S^3} S_{nl}^m = -l(l+2) S_{nl}^m,$$

and form an orthonormal basis defined on the hypersphere, satisfying the normalization condition

$$\int_0^{2\pi} \int_0^\pi \int_0^\pi S_{nl}^m(\beta, \theta, \phi) S_{n'l'}^{m'}(\beta, \theta, \phi) \sin^2 \beta \sin \theta d\beta d\theta d\phi = \delta_{n,n'} \delta_{l,l'} \delta_{m,m'}.$$

Thus, the function r in (6), which maps the unitary hypersphere S^3 to the boundary of a generic domain will be approximated by the expansion

$$(8) \quad r(\beta, \theta, \phi) \approx r_N(\beta, \theta, \phi) = \sum_{n=0}^N \sum_{l=0}^n \sum_{m=-l}^l \alpha_{n,l,m} S_{nl}^m(\beta, \theta, \phi).$$

We define the vector $\mathcal{V} \in \mathbb{R}^P$, where $P = \frac{1}{6}(N+1)(N+2)(2N+3)$, containing all the coefficients $\alpha_{n,l,m}$, and the 4D shape optimization procedure will be performed by searching for optimal vectors \mathcal{V} . In this work we considered $N = 10$, which implies that the vector \mathcal{V} has 506 components.

2.3. Generation of points for the MFS. As mentioned in section 2.1, the MFS approximation (4) satisfies the PDE of the problem, and thus we can focus on the approximation of the boundary condition. We will consider a discrete set of collocation points on the boundary where we will impose the boundary conditions of the problem. Thus, we need to have an algorithm for distributing points on the boundary of a 4D domain with boundary defined by (6). Our approach is to define an almost uniformly distributed set of points on the 4D unitary hypersphere and then map these points to the boundary of the domain taking into account the function r .

We start with a 3D unitary sphere parameterized by

$$(9) \quad \begin{cases} x = \sin(\theta) \cos(\phi), \\ y = \sin(\theta) \sin(\phi), \\ z = \cos(\theta) \end{cases}$$

with $\theta \in [0, \pi]$ and $\phi \in [0, 2\pi[$.

A naive choice could be to consider the points generated with equally spaced angles θ and ϕ but it is well known that this procedure does not produce a uniform distribution of points. Instead, we define an integer number M_C corresponding to the number of collocation points that we would like to place and try to have the same





FIG. 1. Algorithm for distributing points on a 3D sphere.

FIG. 2. Points on the sphere generated by our algorithm with $M_C = 500$ (left) and $M_C = 2000$ (right).

variation locally of the angles θ and ϕ and that the product of these two quantities is approximately equal to the average area of each point, i.e.,

$$(10) \quad \Delta\theta \approx \Delta\phi, \quad \Delta\theta\Delta\phi \approx \frac{4\pi}{M_C} \Rightarrow \Delta\theta \approx \Delta\phi \approx \sqrt{\frac{4\pi}{M_C}}.$$

Next, we fix a meridian, for example corresponding to $\phi = 0$ whose length is π . Thus, in order to have (10), the number of divisions along this meridian is equal to $\lceil \sqrt{\frac{M_C\pi}{4}} \rceil$ (Figure 1(left)). Finally, each of the divisions on the meridian corresponds to a parallel defined by $\theta = \theta_i$. This parallel is a circumference of radius $\sin(\theta_i)$ and, thus has perimeter equal to $2\pi \sin(\theta_i)$. Again, in order to have (12) we must place $\lceil \sqrt{\frac{M_C\pi}{4}} \sin(\theta_i) \rceil$ points uniformly distributed on the parallel (Figure 1(right)). In Figure 2 we plot the points generated by our algorithm with $M_C = 500$ (left) and $M_C = 2000$ (right). The extension of this algorithm for the 4D sphere is straightforward. We take the 4D unitary sphere,

$$(11) \quad \begin{cases} x = \sin(\beta) \sin(\theta) \cos(\phi), \\ y = \sin(\beta) \sin(\theta) \sin(\phi), \\ z = \sin(\beta) \cos(\theta), \\ w = \cos(\beta) \end{cases}$$

with $\beta \in [0, \pi]$, $\theta \in [0, \pi]$, and $\phi \in [0, 2\pi[$. Again, we try to have the same variation of the three angles locally and that the product of these two quantities is approximately

equal to the average area of each point, using the fact that the surface area of a 4D unitary sphere is equal to $2\pi^2$,

$$(12) \quad \Delta\theta \approx \Delta\phi \approx \Delta\beta, \quad \Delta\theta\Delta\phi\Delta\beta \approx \frac{2\pi^2}{M_C} \Rightarrow \Delta\theta \approx \Delta\phi \approx \Delta\beta \approx \sqrt[3]{\frac{2\pi^2}{M_C}} := C.$$

Now, the number of 4D *parallels* corresponding to division along the angle β is equal to $\lceil \frac{\pi}{C} \rceil$, and each of these 4D parallels defined by $\beta = \beta_i$ are 3D spheres of radius $\sin(\beta_i)$. Therefore, we can apply the previous algorithm for 3D spheres. The surface area of a 3D sphere of radius R is given by $4\pi R^2$, and, thus, we use the previous algorithm to place $\lceil \frac{4\pi \sin^2(\beta_i)}{C} \rceil$ points on each of the 4D parallels.

This distribution of points on the sphere is not optimal and other possibilities could be considered. Indeed, the problem of distributing N points on the sphere in such a way that we maximize the minimum distance between any pair of points is a classical problem known as Tammes problem [40, 14]. The problem was already solved for some particular small numbers N (e.g., [20, 16, 33]). Another possible criterion for determining optimal locations for the nodes is to calculate the Fekete points x_i , $i = 1, \dots, N$, that minimize the energy

$$E(N) = \sum_{1 \leq i < j \leq N} \frac{1}{|x_i - x_j|}.$$

In this case, the points x_i correspond to charged particles which repel each other according to Coulomb's law. Several works addressed the numerical solution of this problem for an arbitrary number N , even for a high-dimensional hypersphere [18, 22, 41]. Other approaches for the distribution of nodes could be considered. For example, in [7] an algorithm was described for distributing nodes on the boundary of 3D shapes. However, from the computational point of view, all these approaches are more expensive than the algorithm that was described, which allows us to obtain good distributions of an arbitrary number of nodes in less than one second.

In the context of the application of MFS, the most important issue for having highly accurate results is the choice for the location of the source points y_j (e.g., [4, 3, 9, 7]). In this work we will follow the choice proposed in [4, 7]. We take M_C collocation points x_i , $i = 1, \dots, M_C$, on the boundary of the domain and for each of these points we calculate the outward unitary vector n_i , which is normal to the boundary at x_i . The source points are defined by

$$y_i = x_i + \delta n_i,$$

where δ is a parameter chosen such that the source points remain outside $\bar{\Omega}$. The numerical results that we will present in section 3 were obtained with $M_C \approx 7000$ and $\delta \approx 0.2$.

2.4. Eigenvalue calculation. The MFS approximation satisfies the PDE of the eigenvalue problem and thus the approximation for an eigenvalue is determined searching for the values λ such that there exists a (nonzero) MFS function \tilde{u} fitting the null boundary conditions. We used the Betcke–Trefethen subspace angle approach for the eigenvalue calculation (cf. [11]). We distribute randomly some points $z_i \in \Omega$, $i=1, \dots, M_I$, define the matrices $\mathbf{A}_B(\lambda)$, $\mathbf{A}_I(\lambda)$, where

$$[\mathbf{A}_B(\lambda)]_{i,j} = \Phi_\lambda(x_i - y_j), \quad [\mathbf{A}_I(\lambda)]_{i,j} = \Phi_\lambda(z_i - y_j),$$




FIG. 3. Plot of $\sigma(\lambda)$ for the 4D ball with unit volume.

and

$$\mathbf{A}(\lambda) := \begin{bmatrix} \mathbf{A}_B(\lambda) \\ \mathbf{A}_I(\lambda) \end{bmatrix}$$

and calculate the **QR** decomposition of the matrix $\mathbf{A}(\lambda)$,

$$\begin{bmatrix} \mathbf{A}_B(\lambda) \\ \mathbf{A}_I(\lambda) \end{bmatrix} = \begin{bmatrix} \mathbf{Q}_B(\lambda) \\ \mathbf{Q}_I(\lambda) \end{bmatrix} \mathbf{R}(\lambda).$$

Then, we study the evolution of the eigenvalue of $\mathbf{Q}_B(\lambda)$ with smallest magnitude, which will be called $\sigma(\lambda)$. The approximations for the eigenvalues of the Dirichlet Laplacian are the local minima of $\sigma(\lambda)$ (cf. [11]). In Figure 3 we plot $\sigma(\lambda)$ for $\lambda \in [5, 10]$ obtained for the 4D ball with unit volume and we can locate three eigenvalues in this interval. The method is not very sensitive to the choice of interior points $M_{\mathcal{I}}$ and in all the numerical simulations we fixed $M_{\mathcal{I}} = 50$. In Figure 4 we show some results for the convergence to the first eigenvalue, as a function of the number of collocation points $M_{\mathcal{C}}$, and three different choices of δ . We can observe that the method has fast convergence and with $M_{\mathcal{C}} = 3500$ we obtain relative errors close to machine level precision. Moreover, the best results are obtained for larger parameter δ , which typically happens for the ball (e.g., [4, 9, 7]).

2.5. Optimization algorithm. A very useful mathematical tool for solving shape optimization problems involving Dirichlet eigenvalues of the Laplacian is the Hadamard formula of derivation with respect to the domain (e.g., [26]). Consider an application $\Psi(t)$ such that $\Psi : t \in [0, T[\rightarrow W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$ is differentiable at 0 with $\Psi(0) = I$, $\Psi'(0) = V$, where $W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^d)$ is the set of bounded Lipschitz maps from \mathbb{R}^d into itself, I is the identity, and V is a deformation field. Following the notation of [26], we will denote by $\Omega_t = \Psi(t)(\Omega)$, $\lambda_k(t) = \lambda_k(\Omega_t)$, and by u_k an associated normalized eigenfunction in $H_0^1(\Omega)$. The domains considered are smooth (see section 2.2), thus, if we assume that the eigenvalue $\lambda_k(\Omega)$ is simple, then

$$(13) \quad \lambda'_k(0) = - \int_{\partial\Omega} |\nabla u_k|^2 V \cdot n d\sigma.$$




FIG. 4. Convergence results for the first eigenvalue of the 4D ball.

We also know the derivative of the volume. Define the function $Vol(t) = |\Omega_t|$, then, we have

$$(14) \quad Vol'(0) = \int_{\partial\Omega} V \cdot n \, d\sigma.$$

Now we note that we know that given an homothety of ratio t , we have

$$\lambda_k(t\Omega) = \frac{\lambda_k(\Omega)}{t^2}$$

and

$$|t\Omega| = t^4 |\Omega|,$$

thus, instead of the optimization problem (2), we can study an equivalent problem which avoids the geometrical constraint

$$(15) \quad \lambda_k^* = \text{Min} \left\{ \lambda_k(\Omega) |\Omega|^{\frac{1}{2}} : \Omega \subset \mathbb{R}^4 \right\}, \quad k = 1, 2, \dots$$

It is well known that the minimizers typically correspond to domains for which the optimal eigenvalue is multiple and we must deal with the nonsmoothness of the cost function,

$$\mathcal{F}_k(\mathcal{V}) = \lambda_k(\Omega) |\Omega|^{\frac{1}{2}},$$

where Ω is obtained from \mathcal{V} , by using (6) and (8). Note that the directional derivatives of the eigenvalues with respect to a given perturbation field exist even in the case of multiple eigenvalues (e.g., [26]). We will denote by $g_k \in \mathbb{R}^P$ the shape gradient of \mathcal{F}_k , that is a vector whose i th component is the shape derivative of \mathcal{F}_k with respect to the i th component of the vector \mathcal{V} and split the optimization procedure into two steps. At the first stage, we apply the *quasi*-Newton method LBFGS to the minimization of \mathcal{F}_k (e.g., [30]). Once we are close to a multiple eigenvalue, that is $\mathcal{F}_k - \mathcal{F}_{k-1}$ is small enough, we switch the way we calculate a search direction for performing a line search. For a given (unitary) vector $v \in \mathbb{R}^P$ and small t we have

$$\mathcal{F}_k(\mathcal{V} + tv) \approx \mathcal{F}_k(\mathcal{V}) + t(g_k \cdot v),$$

where $(g_k.v)$ is the directional derivative of \mathcal{F}_k on the direction defined by v . In a similar fashion we have

$$\mathcal{F}_{k-1}(\mathcal{V} + tv) \approx \mathcal{F}_{k-1}(\mathcal{V}) + t(g_{k-1}.v).$$

We want to decrease both \mathcal{F}_k and \mathcal{F}_{k-1} , thus, v must be such that $(g_k.v)$ and $(g_{k-1}.v)$ are both negative and the corresponding absolute values are as large as possible. Thus, it is natural to define the search direction v^* as the solution of the optimization problem

$$(16) \quad \min_{v \in \mathbb{R}^P: \|v\|=1} \max(g_{k-1}.v, g_k.v).$$

The generalization for an eigenvalue with multiplicity m is straightforward and instead of (16), the optimal search direction v^* solves

$$(17) \quad m^* = \min_{v \in \mathbb{R}^P: \|v\|=1} \max(g_{k-m+1}.v, g_{k-m+2}.v, \dots, g_k.v).$$

Another approach would be to use the formula for the derivative of a multiple eigenvalue (e.g., [23, 39]). The optimization procedure is summarized in Algorithm 1.

3. Numerical results. Analyzing geometrical properties of 4D shapes requires special attention. In this spectral theory context, we focus on the identification of symmetries. Actually, the question of the existence of one or many symmetries of optimal profiles minimizing eigenmodes is one of the crucial issues in spectral optimization. This interest is motivated by the fact that symmetries are strongly connected to the multiplicity of eigenvalues in a nontrivial way. We describe below a simple numerical approach that we developed to deduce symmetries of computed optimal shapes by Algorithm 1. The following postprocessing is completely independent of previous optimization and could be used in a different context to identify orthogonal symmetries.

Due to the spherical parameterization (6), all the obtained profiles are star shaped. This fact makes the identification of symmetries easier: consider a sampling of the unit sphere S^3 associated with a finite collection of parameters $(\beta_i, \theta_i, \phi_i)$. For every optimal domain Ω , we computed an analytic radius function $r_\Omega(\beta, \theta, \phi)$ which associates with $(\beta_i, \theta_i, \phi_i)$ by (6) a list of points (P_i) of $\partial\Omega$. Actually, since Ω is star shaped, this map X_Ω from S^3 to $\partial\Omega$ defined by the system (6) is one to one. Consider now s_H , the symmetric transformation with respect to an hyperplane H . That is, $s_H(x) = x + 2 * (p_H(x) - x)$, where p_H stands for the orthogonal projection on H . If H is a hyperplane of symmetry of Ω , we must have

$$(18) \quad \|s_H(X_\Omega(\beta, \theta, \phi))\| = r_\Omega(X_\Omega^{-1}(s_H(X_\Omega(\beta, \theta, \phi)))).$$

We used the previous equality to identify numerically the hyperplane of symmetries of a given shape. Actually, we define a simple cost function on the set of hyperplanes of \mathbb{R}^4 , associated with the sampling $(\beta_i, \theta_i, \phi_i)$, by

$$(19) \quad F(H) = \sum_i \left[\|s_H(X_\Omega(\beta_i, \theta_i, \phi_i))\| - r_\Omega(X_\Omega^{-1}(s_H(X_\Omega(\beta_i, \theta_i, \phi_i)))) \right]^2.$$

We parameterize the space of hyperplanes by their implicit equations. Thus, every vector $\nu \in [-1, 1]^5 \setminus \{0\}$ represents a hyperplane and two such vectors represent the

Algorithm 1. Optimization procedure.

```

1: Require  $\mathcal{V}$ ,  $tol_1 \in (0, 1)$ ,  $tol_2 \in (0, 1)$ ,  $i_{max} \in \mathbb{N}$ ,  $j_{max} \in \mathbb{N}$ 
2:  $i \leftarrow 0$ 
3: Calculate  $\mathcal{F}_k$  and  $g_k$ 
4: Repeat
5:    $i \leftarrow i - 1$ .
6:   Do a step of the LBFGS algorithm, do a linesearch, and update  $\mathcal{V}$  and  $\mathcal{F}_k$ 
7:   Calculate  $g_k$ 
8:   if  $\|g_k\| < tol_1$ 
9:     then stop
10:  else
11:    Calculate  $\mathcal{F}_{k-1}$ 
12:    if  $\mathcal{F}_k - \mathcal{F}_{k-1} < tol_2$ 
13:      then go to step 2
14:    end if
15:  end if
16: Until  $i < i_{max}$ 
17:
18: step 2:
19:  $j \leftarrow 0$ 
20:  $m \leftarrow 2$ 
21: Calculate  $g_{k-1}$ 
22: Repeat
23:    $\mathcal{F}^{prev} \leftarrow \mathcal{F}_k$ .
24:    $j \leftarrow j - 1$ .
25:   Calculate a search direction  $v^*$  by solving (17)
26:   Do a linesearch and update  $\mathcal{V}$  and  $\mathcal{F}_k$ 
27:   Calculate  $\mathcal{F}_{k-1}, \dots, \mathcal{F}_{k-m+1}$ , and  $\mathcal{F}_{k-m}$ 
28:   if  $\mathcal{F}_{k-m+1} - \mathcal{F}_{k-m} < tol_2$ 
29:     then  $m \leftarrow m - 1$ .
30:   end if
31:   Calculate  $g_k, g_{k-1}, \dots, g_{k-m+1}$ 
32: Until  $j < j_{max}$  or  $\mathcal{F}_k - \mathcal{F}^{prev} < tol_1$ 

```

same hyperplane if and only if their coordinates are proportional. Notice that problem (19) is of course not convex but has a few parameters. Thus, it is possible to obtain a reasonably good approximation of an optimal H by a global direct search algorithm which only requires cost function evaluations. In the four test cases under consideration below, it has been possible to obtain a first hyperplane H which is an approximated symmetric hyperplane for the set Ω up to a precision of 0.01 for 1000 fixed sampling parameters $(\beta_i, \theta_i, \phi_i)$. This means that every term in the sum (19) is smaller than 0.01. Once a first hyperplane H_1 (described by ν_1) of symmetry has been identified, we tried to complete ν_1 into an orthonormal base for which every hyperplane of coordinates is as close as possible to a hyperplane of symmetry. To that purpose, we consider a second optimization problem which is the minimization of F among vectors of $[-1, 1]^5 \setminus \{0\}$ which are orthogonal to ν_1 . A normalization of that vector defines the second vector of our base. By induction, we define in such a way an orthogonal basis which has been used to obtain the pictures of Figures 5 and 6.




FIG. 5. Orthogonal cuts of the optimal shapes minimizing the third (first row of images) and the fourth (second row of images) Dirichlet eigenvalues.




FIG. 6. Orthogonal cuts of the optimal shapes minimizing the seventh (first row of images) and eighth (second row of images) Dirichlet eigenvalues.

TABLE 1
Optimal values for λ_i^ and the corresponding multiplicity.*

i	Multiplicity	λ_i^*	$\lambda_i(\mathcal{B})$
3	2	53.95	56.50
4	3	57.06	58.59
5	4	58.59	58.59
6	5	67.06	67.06
7	4	76.28	74.57
8	4	79.17	81.39




FIG. 7. *Computed optimal shape associated the third (left) and the sixth eigenvalue (right) of the Dirichlet Laplacian in dimension 3 under measure constraint.*

More precisely, we first rotated every computed shape with respect to its associated orthogonal basis. In a second step, we estimated the width of the shape in the direction given by the first vector of that basis. Since we always found at least one axis of symmetry in our experiments, this first basis direction is always an axis of symmetry for the computed optimal shape. Finally, we sampled uniformly the width in every four direction of the basis to obtain the orthogonal cuts given in Figures 5 and 6. Every row corresponds to one direction.

4. Discussion of our numerical results. The minimizers of λ_1 and λ_2 are known to be (respectively) a ball and two balls of the same volume (cf. [19, 28, 38, 29]). We considered the minimization of λ_i , for $i = 3, 4, \dots, 8$. The results are summarized in Table 1. The last column shows the best results obtained for unions of balls. The results presented in Table 1 show that our numerical optimizer for λ_3 has a smaller third eigenvalue than the union of three identical balls, which is the best union of balls for this eigenvalue. These numerical estimates provide a counterexample to the Conjecture “Open Problem 8” in [26]. Our results also suggest that the minimizer for λ_4 is a nontrivial connected domain, while the fifth eigenvalue is minimized by the ball and the sixth is minimized by a union of two balls of different volumes, in such a way that the first eigenvalue of the smallest ball is equal to the fifth eigenvalue of the largest ball. To conclude this numerical study, let us point out a striking fact: as in the 3D case, cuts of optimal profile in four dimensions display a strong similarity with optimal domains in lower dimensions. For instance, the computed optimal shape for λ_3 in 4D has cuts (see Figure 5) which look very close to the optimal shape which is suspected to minimize λ_3 under volume constraint in three dimensions (see the left picture in Figure 7). Analogously, some cuts of the computed optimal shape for λ_8 in 4D have a profile very similar to the 3D optimal shape associated with the sixth

eigenvalue (see the right picture in figure 7). We think that a better understanding of these dimensional correlations would lead to deep and new results in the area.

REFERENCES

- [1] V. AKÇELİK, L.-Q. LEE, Z. LI, C. NG, L. XIAO, AND K. KO, *Large scale shape optimization for accelerator cavities*, J. Phys. Conf. Ser., 180 (2009), 012001.
- [2] G. ALLAIRE, S. AUBRY, AND F. JOUVE, *Eigenfrequency optimization in optimal design*, Comput. Methods Appl. Mech. Engrg., 190 (2001), pp. 3565–3579.
- [3] C. J. S. ALVES, *On the choice of source points in the method of fundamental solutions*, Eng. Anal. Bound. Elem., 33 (2009), pp. 1348–1361.
- [4] C. J. S. ALVES AND P. R. S. ANTUNES, *The method of fundamental solutions applied to the calculation of eigenfrequencies and eigenmodes of 2D simply connected shapes*, Comput. Mater. Continua, 2 (2005), pp. 251–266.
- [5] C. J. S. ALVES AND P. R. S. ANTUNES, *The method of fundamental solutions applied to some inverse eigenproblems*, SIAM J. Sci. Comput., 35 (2013), pp. A1689–A1708.
- [6] P. R. S. ANTUNES AND P. FREITAS, *Numerical optimization of low eigenvalues of the Dirichlet and Neumann Laplacians*, J. Opt. Theor. Appl., 154 (2012), pp. 235–257.
- [7] P. R. S. ANTUNES, *Numerical calculation of eigensolutions of 3D shapes using the method of fundamental solutions*, Numer. Methods Partial Differential Equations, 27 (2011), pp. 1525–1550.
- [8] P. R. S. ANTUNES, *Is it possible to tune a drum?*, J. Comp. Phys. 338 (2017), pp. 91–106.
- [9] A. H. BARNETT AND T. BETCKE, *Stability and convergence of the method of fundamental solutions for Helmholtz problems on analytic domains*, J. Comput. Phys., 227 (2008), pp. 7003–7026.
- [10] E. BERTCHIO, A. FERRERO, AND F. GAZZOLA, *Structural instability of nonlinear plates modelling suspension bridges: Mathematical answers to some long-standing questions*, Nonlinear Anal. Real World Appl., 28 (2016), pp. 91–125.
- [11] T. BETCKE AND L. N. TREFETHEN, *Reviving the method of particular solutions*, SIAM Rev., 47 (2005), pp. 469–491.
- [12] D. BUCUR, *Minimization of the k -th eigenvalue of the Dirichlet Laplacian*, Arch. Ration. Mech. Anal., 206 (2012), pp. 1073–1083.
- [13] J. T. CHEN, I. L. CHEN, AND Y. T. LEE, *Eigensolutions of multiply connected membranes using the method of fundamental solutions*, Eng. Anal. Bound. Elem., 29 (2005), pp. 166–174.
- [14] J. H. CONWAY AND N. J. A. SLOANE, *Sphere Packings, Lattices and Groups*, 2nd ed., Springer, New York, 1993.
- [15] S. J. COX AND P. X. UHLIG, *Where best to hold a drum fast*, SIAM Rev., 45 (2003), pp. 75–92.
- [16] L. DANZER, *Finite point-sets on S^2 with minimum distance as large as possible*, Discrete. Math., 60 (1986), pp. 3–66.
- [17] S. K. DONALDSON, *Self-dual connections and the topology of smooth 4-manifolds*, Bull. Amer. Math. Soc., 8 (1983), pp. 81–83.
- [18] T. ERBER AND G. M. HOCKNEY, *Equilibrium configurations of N equal charges on a sphere*, J. Phys. A, 24 (1991), pp. L1369–L1377.
- [19] G. FABER, *Beweis, dass unter allen homogenen membranen von gleicher fläche und gleicher spannung die kreisförmige den tiefsten grundton gibt*, Sitz. ber. bayer. Akad. Wiss. (1923), pp. 169–172.
- [20] L. FEJES TÓTH, *Über die Abschätzung des kürzesten Abstandes zweier Punkte eines auf einer Kugelfläche liegenden Punktsystems*, Jahresber. Dtsch. Math.-Ver 53 (1943), pp. 66–68.
- [21] M. H. FREEDMAN, *The topology of four-dimensional manifolds*, J. Differential Geom., 17 (1982), pp. 357–453.
- [22] L. GLASSER AND A. G. EVERY, *Energies and spacings of point charges on a sphere*, J. Phys. A, 25 (1992), pp. 2473–2482.
- [23] F. DE GOURNAY, *Velocity extension for the level-set method and multiple eigenvalues in shape optimization*, SIAM J. Control Optim., 45 (2006), pp. 343–367.
- [24] E. HABER, *A multilevel level-set method for optimizing eigenvalues in shape design problems*, J. Comp. Phys., 198 (2004), pp. 518–34.
- [25] L. HENRIQUE, J. ANTUNES, AND J. S. CARVALHO, *Shape optimization techniques for musical instrument design*, J. Acoust. Soc. Amer., 110 (2001) 2648.
- [26] A. HENROT, *Extremum Problems for Eigenvalues of Elliptic Operators*, Front. Math., Birkhäuser, Basel, 2006.

- [27] M. KATSURADA, *A mathematical study of the charge simulation method II*, J. Fac. Sci. Univ. Tokyo Sect. IA Math., 36 (1989), pp. 135–162.
- [28] E. KRAHN, *Über eine von Rayleigh formulierte minimaleigenschaft des kreises*, Math. Ann., 94 (1924), pp. 97–100.
- [29] E. KRAHN, *Über Minimaleigenschaften der Kugel in drei und mehr Dimensionen*, Acta Comm. Univ. Dorpat., A9 (1926), pp. 1–44.
- [30] D. C. LIU AND J. NOCEDAL, *On the limited memory method for large scale optimization*, Math. Program. B., 45 (1989), pp. 503–528.
- [31] D. MAZZOLENI AND A. PRATELLI, *Existence of minimizers for spectral problems*, J. Math. Pures App., 100 (2013), pp. 433–453.
- [32] P. M. MORSE AND H. FESHBACH, *Methods of Theoretical Physics, Part I.*, McGraw-Hill, New York, 1953, pp. 623–624.
- [33] O. R. MUSIN AND A. S. TARASOV, *The Tammes Problem for $N = 14$* , Exp. Math., 24 (2015), pp. 460–468.
- [34] D. NORELAND, R. UDAWALPOLA, AND M. BERGGREN, *A hybrid scheme for bore design optimization of a brass instrument*, J. Acoust. Soc. Amer., 128 (2010), pp. 1391–1400.
- [35] E. OUDET, *Numerical minimization of eigenmodes of a membrane with respect to the domain*, ESAIM Control Optim. Calc. Var., 10 (2004), pp. 315–330.
- [36] S. J. OSHER AND F. SANTOSA, *Level set methods for optimization problems involving geometry and constraints. 1: Frequencies of a two-density inhomogeneous drum*, J. Comput. Phys., 171 (2001), pp. 272–288.
- [37] B. OSTING AND C.-Y. KAO, *Minimal convex combinations of sequential Laplace–Dirichlet eigenvalues*, SIAM J. Sci. Comput, 35 (2013), pp. B731–B750.
- [38] G. PÓLYA, *On the characteristic frequencies of a symmetric membrane*, Math. Zeit., 63 (1955), pp. 331–337.
- [39] B. ROUSSELET AND D. CHENAIS, *Continuité et différentiabilité d’éléments propres: Application à l’optimisation de structures*, Appl. Math. Optim., 22 (1990), pp. 27–59.
- [40] R. M. L. TAMMES, *On the origin number and arrangement of the places of exits on the surface of pollengrains*, Rec. Trv. Bot. Neerl., 27 (1930), pp. 1–84.
- [41] N. J. A. SLOANE, *Spherical codes: Nice Arrangements of points on a Sphere in Various Dimensions*, <http://neilsloane.com/packings/index.html>.
- [42] C. H. TAUBES, *Gauge theory on asymptotically periodic 4-manifolds*, J. Differential Geom., 25 (1987), pp. 363–430.
- [43] S. A. WOLF AND J. B. KELLER, *Range of the first two eigenvalues of the Laplacian*, R. Soc. Lond. Proc. Ser. A, Math. Phys. Sci. Eng., 447 (1994), pp. 397–412.